

Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interaction. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

Given the importance of membrane bound proteins, efforts are under way to identify novel membrane bound proteins. Moreover, given the importance of disulfide bond-forming enzymes and their potential uses in a number of different applications, for example in increasing the yield of correct refolding of recombinantly produced proteins, efforts are currently being undertaken by both industry and academia to identify new, native proteins having sequence identity with protein disulfide isomerase. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel protein disulfide isomerase homologs. We herein describe a novel polypeptide having sequence identity with protein disulfide isomerase and the nucleic acids encoding the same.

64. PRO697

Secreted frizzled related proteins (sFRPs) are related to the frizzled family of transmembrane receptors. The sFRPs are approximately 30 kDa in size, and each contains a putative signal sequence, a frizzled-like cysteine-rich domain, and a conserved hydrophilic carboxy-terminal domain. It has been reported that sFRPs may function to modulate Wnt signaling, or function as ligands for certain receptors. Rattner, et al., PNAS USA, 94(7):2859-2863 (1997). Therefore, sFRPs and proteins having sequence identity and/or similarity to sFRPs are of interest.

Another secreted protein of interest is any member of the family of secreted apoptosis-related proteins (SARPs). Expression of SARPs modifies the intracellular levels of beta-catenin, suggesting that SARPs interfere with the Wnt-frizzled proteins signaling pathway. Melkonyan, et al., PNAS USA, 94(25):13636-13641 (1997). Therefore, SARPs and proteins having sequence identity and/or similarity to SARPs are of interest.

In addition to sFRPs and SARPs, many extracellular proteins are of interest. Extracellular proteins play an important role in the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

Secreted proteins have various industrial applications, including pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents.

Efforts are being undertaken by both industry and academia to identify new, native secreted proteins, particularly those having sequence identity or similarity with sFRP-2 and SARP-1. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted

proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., Proc. Natl. Acad. Sci., 93:7108-7113 (1996); U.S. Patent No. 5,536,637)].

5 65. **PRO717**

Efforts are being undertaken by both industry and academia to identify new, native transmembrane receptor proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor proteins. The results of such efforts are provided herein.

10 66. **PRO731**

Cadherins are a large family of transmembrane proteins. Cadherins comprise a family of calcium-dependent glycoproteins that function in mediating cell-cell adhesion in virtually all solid tissues of multicellular organisms. At least cadherins 1-13 as well as types B, E, EP, M, N, P and R have been characterized. Among the functions cadherins are known for, with some exceptions, cadherins participate in cell aggregation and are associated with cell-cell adhesion sites. Recently, it has been reported that while all cadherins share multiple repeats of a cadherin specific motif believed to correspond to folding of extracellular domains, members of the cadherin superfamily have divergent structures and, possibly, functions. In particular it has been reported that members of the cadherin superfamily are involved in signal transduction. See, Suzuki, J. Cell Biochem., 61(4):531-542 (1996). Cadherins are further described in Tanihara, et al., J. Cell Sci., 107(6):1697-1704 (1994), Aberle, et al., J. Cell Biochem., 61(4):514-523 (1996) and Tanihara, et al., Cell Adhes. Commun., 2(1):15-26 (1994).

Protocadherins are members of the cadherin superfamily which are highly expressed in the brain. In some studies, protocadherins have shown cell adhesion activity. See, Sano, et al., EMBO J., 12(6):2249-2256 (1993). However, studies have also shown that some protocadherins, such as protocadherin 3 (also referred to as Pcdh3 or pc3), do not show strong calcium dependent cell aggregation activity. See, Sago, et al., Genomics, 29(3):631-640 (1995) for this study and further characteristics of Pcdh3.

Therefore, novel members of the cadherin superfamily are of interest. More generally, all membrane-bound proteins and receptors are of interest. Such proteins can play an important role in the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and

nerve growth factor receptor.

Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interaction. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

Efforts are therefore being undertaken by both industry and academia to identify new, native membrane bound proteins, particular those having sequence identity with protocadherins, especially 4, 68, 43, 42, 3 and 5. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel membrane-bound proteins. Provided herein are the results of such efforts.

67. PRO218

Efforts are being undertaken by both industry and academia to identify new, native membrane bound proteins, particularly those having sequence identity with membrane regulator proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor proteins.

68. PRO768

The integrins comprise a supergene family of cell-surface glycoprotein receptors that promote cellular adhesion. Each cell has numerous receptors that define its cell adhesive capabilities. Integrins are involved in a wide variety of interaction between cells and other cells or matrix components. The integrins are of particular importance in regulating movement and function of immune system cells. The platelet IIb/IIIa integrin complex is of particular importance in regulating platelet aggregation. A member of the integrin family, integrin β -6, is expressed on epithelial cells and modulates epithelial inflammation. Another integrin, leucocyte-associated antigen-1 (LFA-1) is important in the adhesion of lymphocytes during an immune response.

Of particular interest is H36-alpha 7, an integrin alpha chain that is developmentally regulated during myogenesis as described in Song, et al., *J. Cell Biol.*, 117(3):643-657 (1992). The expression pattern of the laminin-binding alpha 7 beta 1 integrin is developmentally regulated in skeletal, cardiac, and smooth muscle. Ziober, et al., *Mol. Biol. Cell*, 8(9):1723-1734 (1997). It has been reported that expression of the alpha 7-X1/X2 integrin is a novel mechanism that regulates receptor affinity states in a cell-specific context and may modulate integrin-dependent events during muscle development and repair. *Id.* It has further been reported that laminins promote the locomotion of skeletal myoblasts via the alpha 7 integrin receptor. In particular it was reported that alpha 7 beta 1 receptor can promote myoblast adhesion and motility on a restricted number of laminin isoforms and may be important in myogenic precursor recruitment during regeneration and differentiation. Yao, et al., *J. Cell Sci.*, 109(13):3139-3150 (1996). Spliced variants of integrin alpha 7 are also described in Leung, et al., *Biochem. Biophys. Res. Commun.*, 243(1):317-325 (1998) and Fornaro and Languino, *Matrix Biol.*, 16(4):185-193 (1997). Moreover, it has been reported that absence of integrin alpha 7 causes a form of muscular dystrophy. Thus integrins, particularly those related to integrin 7 and related molecules, are of interest.

In addition to the interest of integrins, more generally, all membrane-bound proteins and receptors are